AREDAND DISSEMINATED BY CENTRAL INTELLIGENCE AGENCY Rungary Pressure Chamber at Military Fliers' Hospital This report is the result of a joint collection effort by the Air Force, the Army, the Navy, and CIA and is disseminated in accordance with the provisions of NSCID #17. A pressure chamber is operated by the Magyar Akademia Orvosi Osztaly Tudomanyos Minosito Bizottsag (Rungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budaest IX. The spherical chamber, about 3 to 32 meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 14 m long, 5 to 6 m high, with a flat roof. is attached to the doctors' longe which in turn is connected by a corridor to the hospital laboratory. The chamber had two seats but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Tibor Halm was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the free house to the chamber for "alapanyag vizsgalat" (basic material testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons ad any verse tested but a maxipum of 40 to 50 could be tested. A doctor accompanied each test group.		Third Divine	PAGE FOR SPECIAL CONTROLS, IF ANY This material contains information affecting the
CENTRAL INTELLIGENCE AGENCY Rungary Pressure Chamber at Military Fliers' Hospital This is investigated by the Magyar Akademia Corocal Ostally Tudomanyos Minosito Bizottasg (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budanest IX. The spherical Chamber, about 3 to 33 meters in diameter, is housed in a consectory yellow wooden prefabricated building at the structure is attached to the doctors' Louge which in turn is connected by a corridor to the hospital liboratory. The chamber had the scate building at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber were first given a physical examination, then were brought to the chamber as sellar in it. The interior of the hamber were scate but image work or such as in the chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alaganyae vissgalati" (basic material testing), and the period of time they could do vithout oxygen was ascertained by the Knoch method. An average of 30 persons a day were tested but a maximum of 40 to 50 could be tested. A doctor accompanied each test group. Line location of the Military Fliers' Hospital in Budapest; also a sketch mum of 40 to 50 could be tested. A doctor accompanied each test group.	D	INFORMATION REPORT	National Defense of the United States within th
This report is the result of a joint collection effort by the Air Force, the Army, the Navy, and CIA and is disseminated in accordance with the provisions of NSCID #ff. A pressure chamber is operated by the Magyar Akademia Oravei Osztaly Tudosanyos Minosito Bizottasg (Rungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budanest IX. The spherical chamber, about 3 to 3½ meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 14 m long, 5 to 6 m high, with a flat roof. Is attached to the doctors' lounge which in turn is connected by a corridor to the hospital laboratory. The chamber had to easts but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Tibor Halm was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alapanyag vizsgalat" (basic material testing), and the period of time they could do vitiout oxygen was ascertained by the Kroch method. A average of 30 persons a day were tested but a maximum of 40 to 50 could be tested. A doctor accompanied each test group. **The substitute of the Military Pliers' Mospital in Budapest; also a sketch showing of the pressure chamber building at this hospital.* - end -	PARE		Secs. 793 and 794, the transmission or revelatio
Pressure Chamber at Military Fliers' Hospital This is linevalifated in collection effort by the Air Force, the Army, the Ravy, and CIA and is disseminated in accordance with the provisions of MSCID #ff. A pressure chamber is operated by the Magyar Akademia Orwosi Osstally Tudomanyos Minosito Bizottsag (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gall ut, Budapest IX. The spherical chamber, about 3 to 3 meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 1½ m long, 5 to 6 m high, with a flat roof. Is attached to the doctors' lounge which in turn is connected by a corridor to the hospital laboratory. The chamber had two seats but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Tibor Halm was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alapanyag vizsgalatt" (basic material testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons a day were tested but a maxitation, then were brought to the chamber for "alapanyag vizsgalatt" (basic material testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons a day were tested but a maxitation, then were brought to the chamber for "alapanyag vizsgalatt" (basic material testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons a day were tested but a maxitation of the pressure chamber building at this hospital.) The location of the Military Fliers' Mospital in Budapest; also a sketch showing the control of the pressure chamber building at this hospital.) The locatio	TRY	CENTRAL INTELLIGENCE AGENCY	
Pressure Chamber at Military Fliers' Hospital This report is the result of a joint collection effort by the Air Force, the Army, the Navy, and CIA and is disseminated in accordance with the provisions of NSOID #1. A pressure chamber is operated by the Magyar Akademia Orvosi Osztaly Tudomanyos Minosito Bizottsag (Hungarian Academy of Sciences - Scientific Examination and Personnal Testing) at 17 Gyali ut. Budanest IX. 25X The spherical chamber, about 3 to 3½ meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 14 m long, 5 to 6 m high, with a flat roof. Built after 1945, the structure is attached to the doctors' lounge which in turn is connected by a corridor to the hospital laboratory. The chamber had two seats but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Thor Hain was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alapanyag vissgalat" (basic material testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons a day were tested but a maxistential restriction of the Pressure chamber building at this hospital. The nospital was pretty well supplied with antibiotics so that detrimental after-effects seldom occurred.	IRY	Hungary	
This report is the result of a joint collection effort by the Air Force, the Army, the Havy, and CIA and is disseminated in accordance with the provisions of NSCID #17. 1. A pressure chamber is operated by the Magyar Akademia Orvosi Osztaly Tudomanyos Minosito Bisottsag (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budanest IX. 25X 25X 25X 25X 25X 25X 25X 2	СТ		DATE DISTRIBUTED
This report is the result of a joint collection effort by the Air Force, the Army, the Havy, and CIA and is disseminated in accordance with the provisions of NSCID #17. 1. A pressure chamber is operated by the Magyar Akademia Orvosi Osztaly Tudomanyos Minosito Bisottsag (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budanest IX. 25X 25X 25X 25X 25X 25X 25X 2		Pressure Chamber at Military Flience Hognital	19 mar 57
This report is the result of a joint collection effort by the Air Force, the Army, the Navy, and CIA and is disseminated in accordance with the provisions of NSCID #17. 25X 1. A pressure chamber is operated by the Magyar Akademia Orvosi Osztaly Tudomanyos Minosito Bizottsag (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut. Budancet IX. 25X 25X 25X 25X 25X 25X 25X 2		and the same and t	
/This report is the result of a joint collection effort by the Air Force, the Army, the Navy, and CIA and is disseminated in accordance with the provisions of NSCID #17. 25X A pressure chamber is operated by the Magyar Akademia Orvosi Osztaly Tudomanyos Minosito Bizottsag (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budapest IX. 25X The spherical chamber, about 3 to 32 meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 14 m long, 5 to 6 m high, with a flat roof. 18 attached to the coctors' lounge which in turn is connected by a corridor to the hospital laboratory. The chamber had two seats but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Tibor Halm was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. 3. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alapanyag vizsgalat" basic material testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons a day were tested but a maxitation of 40 to 50 could be tested. A doctor accompanied each test group. 3. Personnel to say pretty well supplied with antibiotics so that detrimental after-effects seldom occurred. 3. Sketch showing the location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital. 4. Location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital. 5. end. 5. Location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital.			SUPPLEMENT TO REPORT #
/This report is the result of a joint collection effort by the Air Force, the Army, the Navy, and CIA and is disseminated in accordance with the provisions of NSCID #17. 25X A pressure chamber is operated by the Magyar Akademia Orvosi Osztaly Tudomanyos Minosito Bizottsag (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budapest IX. 25X The spherical chamber, about 3 to 32 meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 14 m long, 5 to 6 m high, with a flat roof. 18 attached to the coctors' lounge which in turn is connected by a corridor to the hospital laboratory. The chamber had two seats but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Tibor Halm was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. 3. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alapanyag vizsgalat" basic material testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons a day were tested but a maxitation of 40 to 50 could be tested. A doctor accompanied each test group. 3. Personnel to say pretty well supplied with antibiotics so that detrimental after-effects seldom occurred. 3. Sketch showing the location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital. 4. Location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital. 5. end. 5. Location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital.			
25x A pressure chamber is operated by the Magyar Akademia Orwosi Osztaly Tudomanyos Minosito Bizottsag (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budapest IX. 25x The spherical chamber, about 3 to 3½ meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 14 m long, 5 to 6 m high, with a flat roof. Is attached to the footory: lounge which in turn is connected by a corridor to the hospital laboratory. The chamber had two seats but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Tibor Halm was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alapanyag vizsgalat" (basic material testing), and the period of time they could do without oxygen was ascertained by the Krech method. An average of 30 persons a day were tested but a maximum of 40 to 50 could be tested. A doctor accompanied each test group. The location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital. The location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital.		THIS IS LINEVALUATED INFORMAT	ION
25x A pressure chamber is operated by the Magyar Akademia Orwosi Osztaly Tudomanyos Minosito Bizottsag (Hungarian Academy of Sciences - Scientific Examination and Personnel Testing) at 17 Gyali ut, Budapest IX. 25x The spherical chamber, about 3 to 3½ meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 14 m long, 5 to 6 m high, with a flat roof. Is attached to the footory: lounge which in turn is connected by a corridor to the hospital laboratory. The chamber had two seats but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Tibor Halm was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alapanyag vizsgalat" (basic material testing), and the period of time they could do without oxygen was ascertained by the Krech method. An average of 30 persons a day were tested but a maximum of 40 to 50 could be tested. A doctor accompanied each test group. The location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital. The location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital.		This report is the result of a joint collection	officert has the Ato The
The spherical chamber, about 3 to 3½ meters in diameter, is housed in a one-story yellow wooden prefabricated building 12 to 14 m long, 5 to 6 m high, with a flat roof. is attached to the &octors' lounge which in turn is connected by a corridor to the hospital laboratory. The chamber had two seats but "many more" could occupy it at the same time. Instruments were located outside the chamber as well as in it. The interior of the chamber is an insulated room constructed completely of glass. Dr Tibor Halm was chief of the pressure chamber. One or two doctors and an assistant to handle the oxygen tanks were assigned to the chamber. Personnel to be tested in the chamber were first given a physical examination, then were brought to the chamber for "alapanyag vizsgalat" (basic material testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons a day were tested but a maxitum of 40 to 50 could be tested. A doctor accompanied each test group. The nospital was pretty well supplied with antibiotics so that detrimental after-effects seldom occurred. Sketch showing the location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital. **The location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital.	1.	minosito bizottsag (Hungarian Academy of Science	demia Ormosi Osztalu Tudomanuos
testing), and the period of time they could do without oxygen was ascertained by the Kroch method. An average of 30 persons a day were tested but a maximum of 40 to 50 could be tested. A doctor accompanied each test group. The nospital was pretty well supplied with antibiotics so that detrimental after-effects seldom occurred. sketch showing the location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital.	2,		
the location of the Military Fliers' Hospital in Budapest; also a sketch of the pressure chamber building at this hospital.		with a flat roof. is attached to the coctors' lounge which in turn the hospital laboratory. The chamber had two secure it at the same time. Instruments were lowell as in it. The interior of the chamber is a completely of glass. Dr Tibor Halm was chief of or two doctors and an assistant to handle the care.	liameter, is housed in a 2 to 14 m long, 5 to 6 m high, built after 1945, the structure is connected by a corridor to eats but "many more" could coated outside the chamber as in insulated room constructed
of the pressure chamber building at this hospital.		with a flat roof. is attached to the coctors' lounge which in turn the hospital laboratory. The chamber had two se occupy it at the same time. Instruments were lowell as in it. The interior of the chamber is a completely of glass. Dr Tibor Halm was chief of or two doctors and an assistant to handle the ox the chamber. Personnel to be tested in the chamber were first then were brought to the chamber for "alapanyag testing), and the period of time they could do we by the Kroch method. An average of 30 persons as	diameter, is housed in a 2 to 14 m long, 5 to 6 m high, duilt after 1945, the structure is connected by a corridor to eats but "many more" could deated outside the chamber as an insulated room constructed the pressure chamber. One tygen tanks were assigned to given a physical examination, vizsgalat" (basic material ithout oxygen was ascertained day were tested but a mari
of the pressure chamber building at this hospital.	4.	with a flat roof. is attached to the coctors' lounge which in turn the hospital laboratory. The chamber had two secupy it at the same time. Instruments were lowell as in it. The interior of the chamber is a completely of glass. Dr Tibor Halm was chief of or two doctors and an assistant to handle the oather chamber. Personnel to be tested in the chamber were first then were brought to the chamber for "alapanyag testing), and the period of time they could do we by the Kroch method. An average of 30 persons a mum of 40 to 50 could be tested. A doctor accommum of 40 to 50 could be tested with antib	diameter, is housed in a 2 to 14 m long, 5 to 6 m high, will after 1945, the structure is connected by a corridor to eats but "many more" could coated outside the chamber as in insulated room constructed the pressure chamber. One typen tanks were assigned to given a physical examination, vizsgalat" (basic material ithout oxygen was ascertained day were tested but a maxipanied each test group.
	4.	with a flat roof. is attached to the coctors' lounge which in turn the hospital laboratory. The chamber had two secupy it at the same time. Instruments were lowell as in it. The interior of the chamber is a completely of glass. Dr Tibor Halm was chief of or two doctors and an assistant to handle the oather chamber. Personnel to be tested in the chamber were first then were brought to the chamber for "alapanyag testing), and the period of time they could do we by the Kroch method. An average of 30 persons a mum of 40 to 50 could be tested. A doctor accommum of 40 to 50 could be tested with antib	diameter, is housed in a 2 to 14 m long, 5 to 6 m high, suilt after 1945, the structure is connected by a corridor to eats but "many more" could ceated outside the chamber as in insulated room constructed the pressure chamber. One tygen tanks were assigned to given a physical examination, vizsgalat"(basic material ithout oxygen was ascertained day were tested but a maxi- panied each test group. 25) iotics so that detrimental
TION STATE ARMY NAV NAV TARE TAL	4.	with a flat roof. is attached to the coctors' lounge which in turn the hospital laboratory. The chamber had two secupy it at the same time. Instruments were lowell as in it. The interior of the chamber is a completely of glass. Dr Tibor Halm was chief of or two doctors and an assistant to handle the oather chamber. Personnel to be tested in the chamber were first then were brought to the chamber for "alapanyag testing), and the period of time they could do we by the Kroch method. An average of 30 persons a mum of 40 to 50 could be tested. A doctor accommod the country of the chamber selected with antibafter-effects seldom occurred.	diameter, is housed in a 2 to 14 m long, 5 to 6 m high, suilt after 1945, the structure is connected by a corridor to eats but "many more" could reated outside the chamber as in insulated room constructed the pressure chamber. One typen tanks were assigned to given a physical examination, vizsgalat" (basic material ithout oxygen was ascertained day were tested but a maxipanied each test group. sketch showing in Budapest; also a sketch
AIR AIR	4.	with a flat roof. is attached to the doctors' lounge which in turn the hospital laboratory. The chamber had two secure it at the same time. Instruments were lowell as in it. The interior of the chamber is a completely of glass. Dr Tibor Halm was chief of or two doctors and an assistant to handle the oather chamber. Personnel to be tested in the chamber were first then were brought to the chamber for "alapanyag testing), and the period of time they could do we by the Kroch method. An average of 30 persons a mum of 40 to 50 could be tested. A doctor accommum of 40 to 50 could be tested. A doctor accommum of 40 to 50 could be tested. The pressure chamber of the pressure	diameter, is housed in a 2 to 14 m long, 5 to 6 m high, suilt after 1945, the structure is connected by a corridor to eats but "many more" could reated outside the chamber as in insulated room constructed the pressure chamber. One typen tanks were assigned to given a physical examination, vizsgalat" (basic material ithout oxygen was ascertained day were tested but a maxipanied each test group. sketch showing in Budapest; also a sketch
	4.	with a flat roof. is attached to the doctors' lounge which in turn the hospital laboratory. The chamber had two secure it at the same time. Instruments were lowell as in it. The interior of the chamber is a completely of glass. Dr Tibor Halm was chief of or two doctors and an assistant to handle the oather chamber. Personnel to be tested in the chamber were first then were brought to the chamber for "alapanyag testing), and the period of time they could do we by the Kroch method. An average of 30 persons a mum of 40 to 50 could be tested. A doctor accommod to the chamber was pretty well supplied with antibafter effects seldom occurred. The location of the Military Fliers' Hospital is of the pressure chamber end of the pressure pressure the pressure the pressure that the pressure pressure the pressure	diameter, is housed in a 2 to 14 m long, 5 to 6 m high, suilt after 1945, the structure is connected by a corridor to eats but "many more" could reated outside the chamber as in insulated room constructed the pressure chamber. One typen tanks were assigned to given a physical examination, vizsgalat" (basic material ithout oxygen was ascertained day were tested but a maxipanied each test group. sketch showing in Budapest; also a sketch
		with a flat roof. is attached to the doctors' lounge which in turn the hospital laboratory. The chamber had two secure it at the same time. Instruments were lowell as in it. The interior of the chamber is a completely of glass. Dr Tibor Halm was chief of or two doctors and an assistant to handle the oather chamber. Personnel to be tested in the chamber were first then were brought to the chamber for "alapanyag testing), and the period of time they could do we by the Kroch method. An average of 30 persons a mum of 40 to 50 could be tested. A doctor accommod to the chamber was pretty well supplied with antibafter effects seldom occurred. The location of the Military Fliers' Hospital is of the pressure chamber end of the pressure pressure the pressure the pressure that the pressure pressure the pressure	diameter, is housed in a 2 to 14 m long, 5 to 6 m high, suilt after 1945, the structure is connected by a corridor to eats but "many more" could reated outside the chamber as in insulated room constructed the pressure chamber. One typen tanks were assigned to given a physical examination, vizsgalat" (basic material ithout oxygen was ascertained day were tested but a maxipanied each test group. sketch showing in Budapest; also a sketch

Approved For Release 2008/07/01 : CIA-RDP80T00246A000601010001-2

AF FORM 112—PART II
APPROVED 1 JUNE 1948

CONFIDENTIAL (CLASSIFICATION)

25X1

AIR INTELLIGENCE INFORMATION REPORT

FROM (Agency) PAGES 25X1 afoin 1a1, hq usaf 3 PAGE 05 STADIUM KATONAI REPULO KORHAZ OPERATED BY MAGYAR AKADEMIA ORVOSI OSITALY TUDOMANYOS MINIOSITO BIZOTTSAG. GYALI UT 19° 01 OVERLAY OF THE KATONAI REPÜLÖ KORHÁZ (MILITARY FLIERS 25X1

NOTE: THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE ACT, 50 U. S. C.—
31 AND 32, AS AMENDED. ITS TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW,
IT MAY NOT BE REPRODUCED IN WHOLE OR IN PART, BY OTHER THAN UNITED STATES AIR FORCE AGENCIES, EXCEPT BY PERMISSION OF THE DIRECTOR OF
INTELLIGENCE, USAF.

CONFIDENTIAL
Approved For Release 2008/07/01 : CIA-RDP80T00246A000601010001-2 - GOVERNMENT PRINTING OFFICE


AF FORM 112—PART II APPROVED 1 JUNE 1948



25X1

25X1

AIR INTELLIGENCE INFORMATION REPORT



NOTE: THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE ACT, 50 U. S. C.—
31 AND 32, AS AMENDED. ITS TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW,
IT MAY NOT BE REPRODUCED IN WHOLE OR IN PART, BY OTHER THAN UNITED STATES AIR FORCE AGENCIES, EXCEPT BY PERMISSION OF THE DIRECTOR OF
INTELLIGENCE, USAF.

CONFIDENTIAL

Approved For Release 2008/07/01: CIA-RDP80T00246A000601010001-2